# Probing the nature of phases across the phase transition at finite isospin chemical potential

Rajiv V. Gavai\* Department of Theoretical Physics Tata Institute of Fundamental Research Mumbai, INDIA

\* Work in progress with G. S. Bali & G. Endrődi (Regensburg) and N. Mathur (TIFR)

CPOD 2016, University of Wroclaw, Poland June 4, 2016

# Introduction : Why $\mu_I$ ?

- Protons convert to neutrons and neutrinos via electron capture in the core of neutron stars, leading to high baryon density with considerable  $I_3$ , i.e., finite isospin.
- Using  $\mu_u$  and  $\mu_d$  as light quark chemical potentials, one has  $\mu_B = 3(\mu_u + \mu_d)/2$  and  $\mu_I = (\mu_u \mu_d)/2$
- No sign/phase problem for  $\mu_B=0$  but  $\mu_I
  eq 0$ . (Son-Stephanov PRL '01, Kogut-Sinclair PRD '02)

# Introduction : Why $\mu_I$ ?

- Protons convert to neutrons and neutrinos via electron capture in the core of neutron stars, leading to high baryon density with considerable  $I_3$ , i.e., finite isospin.
- Using  $\mu_u$  and  $\mu_d$  as light quark chemical potentials, one has  $\mu_B = 3(\mu_u + \mu_d)/2$  and  $\mu_I = (\mu_u \mu_d)/2$
- No sign/phase problem for  $\mu_B=0$  but  $\mu_I
  eq 0$ . (Son-Stephanov PRL '01, Kogut-Sinclair PRD '02)
- For QCD with two flavours, quarks are two component spinors, leading to a quark matrix :

$$M = \begin{pmatrix} \mathcal{D}(\mu_I) + m & \lambda \gamma_5 \\ -\lambda \gamma_5 & \mathcal{D}(-\mu_I) + m \end{pmatrix}$$

where  $\lambda$  is introduced as an isospin-breaking term to study SSB in  $\lambda \to 0$  limit.

- The fermion determinant can be shown to be real, and for staggered fermions the usual techniques work to simulate the theory, as shown by Kogut-Sinclair (PRD '02), who also obtained first numerical results for the phase transition.
- Employing staggered fermions on  $8^4$  lattices with a = 0.299(2) fm & lattice quark mass ma = 0.025, corresponding to  $m_{\pi} \simeq 260$  MeV, Endrődi (PRD '15) investigated the phase structure.

- The fermion determinant can be shown to be real, and for staggered fermions the usual techniques work to simulate the theory, as shown by Kogut-Sinclair (PRD '02), who also obtained first numerical results for the phase transition.
- Employing staggered fermions on  $8^4$  lattices with a = 0.299(2) fm & lattice quark mass ma = 0.025, corresponding to  $m_{\pi} \simeq 260$  MeV, Endrődi (PRD '15) investigated the phase structure.
- He computed the chiral condensate, the pion condensate, and the isospin density by using,

$$\langle \bar{\psi}\psi \rangle = \frac{T}{V} \frac{\partial \log Z}{\partial m}, \\ \langle \pi \rangle = \langle \bar{\psi}_u \gamma_5 \psi_d - \bar{\psi}_d \gamma_5 \psi_u \rangle = \frac{T}{V} \frac{\partial \log Z}{\partial \lambda}, \\ \langle n_I \rangle = \frac{T}{V} \frac{\partial \log Z}{\partial \mu_I}$$

and obtained  $a\mu_I^c \simeq 0.2$ .





•  $\lambda \to 0$  extrapolation in yellow, points (linear), line (chiral). • Grey vertical band denotes  $m_{\pi}/2$ .

• Pion condensate & Isospin density become nonzero around  $\mu_I^c\simeq m_\pi/2$ , where chiral condensate vanishes as well.



•  $\lambda \to 0$  extrapolation in yellow, points (linear), line (chiral). • Grey vertical band denotes  $m_{\pi}/2$ .

• Pion condensate & Isospin density become nonzero around  $\mu_I^c \simeq m_\pi/2$ , where chiral condensate vanishes as well. • Polyakov loop displayed in the upper half of the left panel shows deconfinement to occur there as well.

# **Introduction II : Nature of Probe**

- High  $\mu$  phase appears to have restored chiral symmetry and deconfinement. Leading candidate for  $\chi$ SB – topological excitations.
- Successful phenomenology built on Instanton-fermion couplings. (Schafer-Shuryak RMP '98, Diakanov hep-ph/9602375)

# **Introduction II : Nature of Probe**

- High  $\mu$  phase appears to have restored chiral symmetry and deconfinement. Leading candidate for  $\chi$ SB – topological excitations.
- Successful phenomenology built on Instanton-fermion couplings. (Schafer-Shuryak RMP '98, Diakanov hep-ph/9602375)
- LQCD simulations also support it : Instanton-distribution peaks at a radius  $\rho = 0.3$  fm (DeGrand-Hasenfratz PRD '01).



# **Introduction II : Nature of Probe**

- High  $\mu$  phase appears to have restored chiral symmetry and deconfinement. Leading candidate for  $\chi$ SB – topological excitations.
- Successful phenomenology built on Instanton-fermion couplings. (Schafer-Shuryak RMP '98, Diakanov hep-ph/9602375)
- LQCD simulations also support it : Instanton-distribution peaks at a radius  $\rho = 0.3$  fm (DeGrand-Hasenfratz PRD '01).



♠ Note that Overlap Dirac operator, which has *exact* chiral symmetry on the lattice as well as an index theorem, was used for the analysis above.

#### The Overlap Dirac operator spectra has also been used to understand the nature of the high temperature phase.

 $\Diamond$  Number of low eigen modes do get depleted as  $T \uparrow$ . (Edwards-Heller-Kiskis-Narayanan, PRL '99, NPB (PS) '00, PRD '01; Gavai-Gupta-Lacaze, PRD '02)

The Overlap Dirac operator spectra has also been used to understand the nature of the high temperature phase.

 $\Diamond$  Number of low eigen modes do get depleted as  $T \uparrow$ . (Edwards-Heller-Kiskis-Narayanan, PRL '99, NPB (PS) '00, PRD '01; Gavai-Gupta-Lacaze, PRD '02)



 $\diamond$  Furthermore, a gap appears to separate the low modes from others.

Overlap Dirac-Neuberger fermions possess satisfy the following correlator equalities in the chirally symmetric phase :  $C_S(z) = -C_{PS}(z)$  and  $C_V(z) = C_{AV}(z)$ .

Overlap Dirac-Neuberger fermions possess satisfy the following correlator equalities in the chirally symmetric phase :  $C_S(z) = -C_{PS}(z)$  and  $C_V(z) = C_{AV}(z)$ .



♠ Localized zero modes seen for 1.25  $\leq T/T_c \leq 2$  :  $U_A(1)$  restored only gradually up to  $2T_c$ . (G-G-L,PRD 2002)

CPOD 2016, University of Wroclaw, Poland June 4, 2016

Overlap Dirac-Neuberger fermions possess satisfy the following correlator equalities in the chirally symmetric phase :  $C_S(z) = -C_{PS}(z)$  and  $C_V(z) = C_{AV}(z)$ .



 $\heartsuit$  Vector and Axial vector correlators  $\bigstar$  Localized zero modes seen for 1.25equal above  $T_c$  but Pseudoscalar and  $\leq T/T_c \leq 2$ :  $U_A(1)$  restored onlyscalar equal only without zero modes gradually up to  $2T_c$ . (G-G-L,PRD 2002)  $(T = 1.5T_c \text{ above})$ .

# **Our Results**

- We employed the Arnoldi method to extract the eigenvalues of Overlap Dirac operator (defined on dynamical configurations with nonzero  $\mu_I$ ), demanding a residue  $r = ||DX \eta|| \le 10^{-10}$ .
- Extracted  $\sim$ 500 eigenvalues from each configuration.

# **Our Results**

- We employed the Arnoldi method to extract the eigenvalues of Overlap Dirac operator (defined on dynamical configurations with nonzero  $\mu_I$ ), demanding a residue  $r = ||DX \eta|| \le 10^{-10}$ .
- Extracted  $\sim$ 500 eigenvalues from each configuration.
- Used a larger  $24^3 \times 6$  lattice and a Symanzik improved action with 2 stout steps at the physical pion mass.
- $a\mu_I^c = 0.1$  here, which again corresponds to  $\mu_I^c$  being  $m_{\pi}/2$ .
- Computations made at two  $\mu_I$  values, below and above the transition and two different  $\lambda$  the isospin breaking parameter in the quark matrix.

### $\lambda=0.0006$

 $\heartsuit$  We examined Overlap Dirac eigenmodes for  $\mu/\mu_I \simeq 0.5$  and 1.5, corresponding to  $a\mu_I = 0.05$  and 0.15 respectively.



# $\lambda=0.0006$

 $\heartsuit$  We examined Overlap Dirac eigenmodes for  $\mu/\mu_I \simeq 0.5$  and 1.5, corresponding to  $a\mu_I = 0.05$  and 0.15 respectively.





 $\heartsuit$  As expected, the overlap is indeed significant. Alternatively, the surprise is confirmed to be not an illusion.

 $\heartsuit$  Looking at the eigenvalue distribution on a log scale, one can easily identify the zero modes from the gap in the spectrum. Explicit chirality checks needed to confirm their nature.



 $\oint \mu/mu_I = 0.5$  : Zero modes get separated from the others visually.

 $\heartsuit$  Looking at the eigenvalue distribution on a log scale, one can easily identify the zero modes from the gap in the spectrum. Explicit chirality checks needed to confirm their nature.



 $\heartsuit$  Zooming in on the eigenvalue distribution on the log scale to see if the near-zero modes have any difference which was missed.



 $\oint \mu/mu_I = 0.5$ : Nice smooth fall-off is seen.

 $\heartsuit$  Zooming in on the eigenvalue distribution on the log scale to see if the near-zero modes have any difference which was missed.





#### $\heartsuit$ No visible difference in the near-zero mode distributions.,

#### $\lambda=0.0025$

 $\heartsuit$  Again we examined Overlap Dirac eigenmodes for  $\mu/\mu_I \simeq 0.5$  and 1.5, corresponding to  $a\mu_I = 0.05$  and 0.15 respectively.



## $\lambda=0.0025$

 $\heartsuit$  Again we examined Overlap Dirac eigenmodes for  $\mu/\mu_I \simeq 0.5$  and 1.5, corresponding to  $a\mu_I = 0.05$  and 0.15 respectively.



# What about Zero Modes?

Nonzero modes are doubly degenerate for Overlap fermions as a result of the chiral symmetry.

- $\diamond$  Zero modes are *not* degenerate & come with specific chirality, +ve or -ve.
- Act as a direct measure of topology.

# What about Zero Modes?

Nonzero modes are doubly degenerate for Overlap fermions as a result of the chiral symmetry.

- $\diamond$  Zero modes are *not* degenerate & come with specific chirality, +ve or -ve.
- Act as a direct measure of topology.
- For  $T \neq 0$ , Gavai-Gupta-Lacaze (PRD '02) found
- $\begin{array}{ccc} T/T_c & N_{zero} \\ 1.25 & 18 \\ 1.5 & 8 \\ 2.0 & 1 \end{array}$
- A steep fall off is seen. Note  $N_{zero}$  substantial near  $T_c$ .

# What about Zero Modes?

A Nonzero modes are doubly degenerate for Overlap fermions as a result of the chiral symmetry.

- $\diamond$  Zero modes are *not* degenerate & come with specific chirality, +ve or -ve.
- Act as a direct measure of topology.
- For  $T \neq 0$ , Gavai-Gupta-Lacaze• For  $\mu_I \neq 0$ , we find for same (PRD '02) found number of configs (50) :

$T/T_c$	$N_{zero}$			$\mu_I/\mu_I^c$	$N_{zero}^{0.0006}$	$N_{zero}^{0.0025}$	
1.25	18			0.5	426	416	
1.5	8			1.5	451	310	
2.0	1			• No va	riation a	cross $\mu_I$	for $\lambda =$
• A steep fall off is seen. Note $N_{zero} 0.0006$ & a mild one for $\lambda = 0.0025$							
substantial near $T_c$ .			(25% re	(25% reduction)			

# Summary

- We investigated the eigenvalue distribution for chirally exact Overlap Dirac operator for  $\mu_I/\mu_I^c = 0.5 \& 1.5$ , *i. e.*, below and above the isospin phase transition.
- The distribution of zero and near-zero modes is nearly the same for both at  $\lambda = 0.0006$ , with a 25 % reduction in former at  $\lambda = 0.0025$ .

# Summary

- We investigated the eigenvalue distribution for chirally exact Overlap Dirac operator for  $\mu_I/\mu_I^c = 0.5 \& 1.5$ , *i. e.*, below and above the isospin phase transition.
- The distribution of zero and near-zero modes is nearly the same for both at  $\lambda = 0.0006$ , with a 25 % reduction in former at  $\lambda = 0.0025$ .
- This should be contrasted with the earlier  $T \neq 0$  results, where too these modes were present above the transition but decreased sharply as one moved away from the transition.
- Further investigations are going on to pin down the changes in the near-zero modes more quantitatively in an effort to understand the difference in T and  $\mu_I$  directions.

CPOD 2016, University of Wroclaw, Poland June 4, 2016