

Symposium on quantum symmetries, noncommutative geometry and quantum gravity

Dedicated to Jurek Lukierski on his 85th birthday,

24 June 2021, Wroclaw, Poland

# Yang-Baxter sigma models from 4D Chern-Simons theory

Dept. of Phys., Kyoto Univ.

Kentaroh Yoshida



Happy 85th Birthday, Jurek!

## Our interest here

Construct a unified way to describe all of the 2D integrable models

Why is this issue important? (My personal point of view)

In the study of integrable systems, integrable models are discovered suddenly and when a certain amount of them have been obtained, beautiful universal structures behind them are extracted such as Yang-Baxter equation.

Even now, new integrable models are being discovered one after another. But we did not know a method to describe everything from the traditional integrable models to the latest new types of models in a unified manner.

If this is compared to the study of elementary particle physics, the discovery of an integrable model corresponds to that of a new particle, and its unified theory corresponds to finding a unified model of elementary particles (though this theory would be replaced by a larger new theory, subsequently,,).)

## The candidate of the unified theory

### 4D Chern-Simons (CS) theory

[Costello-Yamazaki, 1908.02289]

$$S[A] = \frac{i}{4\pi} \int_{\mathcal{M} \times \mathbb{C}P^1} \omega \wedge CS(A)$$

c.f. Costello-Yamazaki-Witten,  
1709.09993, 1802.01579

$A$  takes a value in Lie algebra  $\mathfrak{g}^{\mathbb{C}}$  of a semi-simple Lie group  $G^{\mathbb{C}}$

$\mathcal{M}$  : a 2D surface with the coordinates  $(\tau, \sigma)$  .  $z$  is a coordinate of  $\mathbb{C}P^1$  .

$$CS(A) \equiv \left\langle A, dA + \frac{2}{3}A \wedge A \right\rangle \quad : \text{Chern-Simons 3-form}$$

$$\omega \equiv \varphi(z)dz \quad : \text{a meromorphic 1-form}$$

This 1-form is closely related to the integrable structure of 2D integrable sigma model (ISM) to be derived.

## The recipe to derive 2D ISMs from 4D CS

1. Prepare a meromorphic 1-form.

The structure of poles and zeros determines the resulting 2D ISM.

2. Take a boundary condition for the gauge field  $A$ .

Possible boundary conditions are governed by the equation of motion.

3. Reduce 4D CS to a 2D system by following a procedure.

There are some reduction methods. Take one of them as you like.

As a result, we see that the resulting 2D system is classically integrable because the associated Lax pair can be constructed along this way.

## The content of my talk

Explain how to derive 2D ISMs from 4D CS by taking a reduction method developed by Delduc-Lacroix-Magro-Vicedo (DLMV)

[Delduc-Lacroix-Magro-Vicedo, 1909.13824]

1) A reduction method by DLMV

2) Concrete examples: 2D principal chiral model  
Yang-Baxter sigma models

A brief summary of my related works

[Fukushima-Sakamoto-KY]

3) Summary and discussion

# 1) A reduction method by DLMV

[Delduc-Lacroix-Magro-Vicedo, 1909.13824]

Our starting point:

$$S[A] = \frac{i}{4\pi} \int_{\mathcal{M} \times \mathbb{C}P^1} \omega \wedge CS(A) , \quad CS(A) \equiv \left\langle A, dA + \frac{2}{3} A \wedge A \right\rangle$$

$$\omega \equiv \varphi(z) dz \quad : \text{ a meromorphic 1-form}$$

This action has an extra gauge symmetry:

$$A \mapsto A + \chi dz$$

Hence the  $z$ -component can always be gauged away:

$$A = A_\sigma d\sigma + A_\tau d\tau + A_{\bar{z}} d\bar{z}$$

Equations of motion:

$$\omega \wedge F(A) = 0 \quad (\text{bulk eom})$$



Species of 2D ISM

$$d\omega \wedge \langle A, \delta A \rangle = 0 \quad (\text{boundary eom})$$



Integrable deformation

**NOTE 1 :** If  $\varphi$  is smooth, the boundary eom is trivially satisfied.

But now 
$$d\omega = \partial_{\bar{z}}\varphi(z) d\bar{z} \wedge dz \quad \text{i.e.,} \quad \partial_{\bar{z}}\frac{1}{z} = 2\pi\delta(z, \bar{z})$$

and hence a delta function may appear if  $\varphi$  has a pole.

**NOTE 2 :** From the bulk eom, the zeros of  $\varphi$  are also important because a derivative of  $A$  may be a distribution, i.e.,  $x \delta(x) = 0$  .

Let us introduce the following notation:

$\mathfrak{p}$  : set of poles of  $\varphi$        $\mathfrak{z}$  : set of zeros of  $\varphi$

**NOTE3:** The boundary eom has the support only on  $\mathcal{M} \times \mathfrak{p} \subset \mathcal{M} \times \mathbb{C}P^1$  .

Indeed, it can be rewritten as

$$\sum_{x \in \mathfrak{p}} \sum_{p \geq 0} (\text{res}_x \xi_x^p \omega) \epsilon^{ij} \frac{1}{p!} \partial_{\xi_x}^p \langle A_i, \delta A_j \rangle |_{\mathcal{M} \times \{x\}} = 0$$

Here the local holomorphic coordinates  $\xi_x$  are defined as

$$\xi_x \equiv z - x \quad (x \in \mathfrak{p} \setminus \{\infty\}), \quad \xi_\infty \equiv 1/z$$

## Lax form

Let us perform a formal gauge transformation:

$$A = -d\hat{g}\hat{g}^{-1} + \hat{g} \mathcal{L} \hat{g}^{-1} \quad \text{a smooth function } \hat{g} : \mathcal{M} \times \mathbb{C}P^1 \rightarrow G^{\mathbb{C}}$$

Then the  $\bar{z}$ -component of  $\mathcal{L}$  can be removed as  $\mathcal{L}_{\bar{z}} = 0$

Then the Lax form is given by

$$\mathcal{L} \equiv \mathcal{L}_\sigma d\sigma + \mathcal{L}_\tau d\tau \quad (\text{to be identified with Lax of 2D ISM})$$

The bulk eom leads to

$$\partial_\tau \mathcal{L}_\sigma - \partial_\sigma \mathcal{L}_\tau + [\mathcal{L}_\tau, \mathcal{L}_\sigma] = 0 \quad \longrightarrow \quad \text{Flatness condition}$$

$$\Downarrow \quad \omega \wedge \partial_{\bar{z}} \mathcal{L} = 0$$

**NOTE:** the set of zeros of  $\varphi$  is that of poles of  $\mathcal{L}$

**FACT**

From this information on the pole structure and the boundary condition for  $A$ , one can determine the explicit form of  $\mathcal{L}$  .

The original 4D CS can be rewritten as

$$S[A] = -\frac{i}{4\pi} \int_{\mathcal{M} \times \mathbb{C}P^1} \omega \wedge d\langle \hat{g}^{-1} d\hat{g}, \mathcal{L} \rangle - \frac{i}{4\pi} \int_{\mathcal{M} \times \mathbb{C}P^1} \omega \wedge I_{\text{WZ}}[\hat{g}]$$

$$I_{\text{WZ}}[u] \equiv \frac{1}{3} \langle u^{-1} du, u^{-1} du \wedge u^{-1} du \rangle$$

To reduce this 4D action to a 2D theory, let us suppose

**the archipelago conditions:**

There exist open disks  $V_x, U_x$  for each  $x \in \mathfrak{p}$  such that  $\{x\} \subset V_x \subset U_x$  and

- i)  $U_x \cap U_y = \emptyset$  if  $x \neq y$  for all  $x, y \in \mathfrak{p}$
- ii)  $\hat{g} = 1$  outside  $\mathcal{M} \times \bigcup_{x \in \mathfrak{p}} U_x$
- iii)  $\hat{g}|_{\mathcal{M} \times U_x}$  depends only on  $\tau, \sigma$  and the radial coordinate  $|\xi_x|$
- iv)  $\hat{g}|_{\mathcal{M} \times V_x}$  depends only on  $\tau, \sigma$ , that is,  $g_x \equiv \hat{g}|_{\mathcal{M} \times V_x} = \hat{g}|_{\mathcal{M} \times \{x\}}$

## Master formula

$$S[\{g_x\}_{x \in \mathfrak{p}}] = \frac{1}{2} \sum_{x \in \mathfrak{p}} \int_{\mathcal{M}} \langle \text{res}_x(\varphi \mathcal{L}), g_x^{-1} dg_x \rangle \\ - \frac{1}{2} \sum_{x \in \mathfrak{p}} (\text{res}_x \omega) \int_{\mathcal{M} \times [0, R_x]} I_{\text{WZ}}[g_x]$$

## Recipe of 2D ISM refined

1. Specify the form of  $\omega$
2. Take a boundary condition of  $A$  at the poles of  $\omega$
3. Fix the form of Lax form  $\mathcal{L}$  with the above information.
4. Finally, evaluate the above master formula.



2D ISM

## 2) Concrete examples

### 1. Principal chiral model with Wess-Zumino (WZ) term

A meromorphic 1-form is

$$\omega = K \frac{1 - z^2}{(z - k)^2} \quad K, k : \text{real constants}$$

The boundary condition of  $A$  at the poles of  $\omega$  is

$$A_i|_k = 0, \quad A_i|_\infty = 0 \quad (i = \tau, \sigma)$$

Lax form:

$$\mathcal{L} = \frac{k - 1}{z - 1} j_+ d\sigma^+ + \frac{k + 1}{z + 1} j_- d\sigma^-$$

$$\sigma^\pm = \frac{1}{2}(\tau \pm \sigma)$$

$$j_\pm = g^{-1} \partial_\pm g$$

2D action:

$$S[g] = \frac{K}{2} \int_{\mathcal{M}} d\sigma \wedge d\tau \langle j_+, j_- \rangle + K k I_{\text{WZ}}[g]$$

## 2. Homogeneous Yang-Baxter sigma model

The 1-form is the same as the previous (but  $k=0$  for simplicity)

$$\omega = K \frac{1 - z^2}{z^2} \quad K : \text{ a real constant}$$

But the boundary condition of  $A$  at the poles of  $\omega$  is replaced by

$$A_i|_0 = -R \partial_z A_i|_0, \quad A_i|_\infty = 0 \quad (i = \tau, \sigma)$$

Here  $R$  is a linear operator from  $\mathfrak{g} \rightarrow \mathfrak{g}$  satisfying

the homogeneous Yang-Baxter equation

$$[R(x), R(y)] - R([R(x), y] + [x, R(y)]) = 0$$

It is useful to introduce the notation:  $R_g \equiv \text{Ad}_{g^{-1}} \circ R \circ \text{Ad}_g$

Lax form: 
$$\mathcal{L} = \frac{1}{z-1} \frac{-1}{1+R_g} j_+ d\sigma^+ + \frac{1}{z+1} \frac{1}{1-R_g} j_- d\sigma^-$$

2D action: 
$$S[g] = \frac{K}{2} \int_{\mathcal{M}} d\sigma \wedge d\tau \left\langle j_+, \frac{1}{1-R_g} j_- \right\rangle$$

### Homogeneous Yang-Baxter sigma model


[Klimcik, hep-th/0210095, 0802.3518] [Delduc-Magro-Vicedo, 1308.3581] [Matsumoto-KY, 1501.03665]

### My related works:

- Homogeneous YB deformed  $\text{AdS}_5 \times \text{S}^5$  superstring [Kawaguchi-Matsumoto-KY, 1401.4855]  
From 4D CS [Fukushima-Sakamoto-KY, 2005.04950]
- Faddeev-Reshetikhin model [Faddeev-Reshetikhin, Ann. Phys. 167 (1986) 227]  
From 4D CS [Fukushima-Sakamoto-KY, 2012.07370]
- Integrable  $T^{1,1}$  sigma model [Arutyunov-Bassi-Lacroix, 2010.05573]  
From 4D CS [Fukushima-Sakamoto-KY, 2105.14920]

[» Mathematics](#) [» Mathematical Physics](#)

[SpringerBriefs in Mathematical Physics](#)



© 2021

## Yang–Baxter Deformation of 2D Non-Linear Sigma Models

Towards Applications to AdS/CFT

Authors: Yoshida, Kentaroh

Introduces a new method called Yang–Baxter deformation to perform integrable deformations systematically

[» see more benefits](#)

### About this book

In mathematical physics, one of the fascinating issues is the study of integrable systems. In particular, non-perturbative techniques that have been developed have triggered significant insight for real physics. There are basically two notions of integrability: classical integrability and quantum integrability. In this book, the focus is on the former, classical integrability. When the system has a finite number of degrees of freedom, it has been well captured by the Arnold–Liouville theorem. However, when the number of degrees of freedom is infinite, as in classical field theories, the

[» Show all](#)

### 3) Summary and Discussion

We have discussed how to derive 2D ISMs from 4D CS.

It is very significant to check if Costello-Yamazaki conjecture is true or not.

#### My previous work with Jurek

Yang-Baxter deformations of Minkowski spacetime with the  $\kappa$ -Poincare  $r$ -matrices.

[Borowiec-Kyono-Lukierski-Sakamoto-KY, 1510.03083]

It may be interesting to consider a connection between 4D CS and our previous work. Flat space limit? Contraction of the algebra?

This issue is my birthday present for Jurek!

*Thank you  
for your  
attention!*